

**Solutions to Practice Test A for Midterm Exam 1**

**1.** [30 Points] Evaluate each of the following limits. Please **justify** your answers. Be clear if the limit equals a value,  $+\infty$  or  $-\infty$ , or Does Not Exist.

(a)  $\lim_{x \rightarrow -3} \frac{x^2 - 2x - 15}{x^2 + x - 6} =$

(b)  $\lim_{x \rightarrow 5} \frac{x^2 - 2x - 15}{|5 - x|} =$

(c)  $\lim_{x \rightarrow 2} \frac{x^2 - 2x - 15}{x^2 + x - 6} =$

(d)  $\lim_{x \rightarrow 5} \frac{x^2 - 2x - 15}{x^2 + x - 6} =$

(e)  $\lim_{x \rightarrow 2} \frac{x + 7}{(x - 2)^2} =$

(f)  $\lim_{x \rightarrow -1} \frac{H(x + 1) - H(-1 - x)}{x + 1} =$  where  $H(x) = \sqrt{x + 2}$

**Solutions.** (a):  $\lim_{x \rightarrow -3} \frac{x^2 - 2x - 15}{x^2 + x - 6} = \lim_{x \rightarrow -3} \frac{(x + 3)(x - 5)}{(x + 3)(x - 2)} = \lim_{x \rightarrow -3} \frac{x - 5}{x - 2} \stackrel{\text{DSP}}{=} \frac{-8}{-5} = \boxed{\frac{8}{5}}$

(b):  $\lim_{x \rightarrow 5} \frac{x^2 - 2x - 15}{|5 - x|}$  is piecewise defined right at  $x = 5$ , so we check both sides:

LHL =  $\lim_{x \rightarrow 5^-} \frac{x^2 - 2x - 15}{|5 - x|} = \lim_{x \rightarrow 5^-} \frac{(x + 3)(x - 5)}{-(x - 5)} = \lim_{x \rightarrow 5^-} -(x + 3) \stackrel{\text{DSP}}{=} -8$

RHL =  $\lim_{x \rightarrow 5^+} \frac{x^2 - 2x - 15}{|5 - x|} = \lim_{x \rightarrow 5^+} \frac{(x + 3)(x - 5)}{(x - 5)} = \lim_{x \rightarrow 5^+} (x + 3) \stackrel{\text{DSP}}{=} 8$

LHL  $\neq$  RHL, so the original limit DNE

(c):  $\lim_{x \rightarrow 2} \frac{x^2 - 2x - 15}{x^2 + x - 6} = \lim_{x \rightarrow 2} \frac{(x + 3)(x - 5)}{(x + 3)(x - 2)} = \lim_{x \rightarrow 2} \frac{x - 5}{x - 2}$  is  $\frac{-3}{0}$ , so we check both sides:

LHL =  $\lim_{x \rightarrow 2^-} \frac{x - 5}{x - 2} = \frac{-3}{0^-} = +\infty$

RHL =  $\lim_{x \rightarrow 2^+} \frac{x - 5}{x - 2} = \frac{-3}{0^+} = -\infty$

LHL  $\neq$  RHL, so the original limit DNE

(d):  $\lim_{x \rightarrow 5} \frac{x^2 - 2x - 15}{x^2 + x - 6} \stackrel{\text{DSP}}{=} \frac{25 - 10 - 15}{25 + 5 - 6} = \frac{0}{24} = \boxed{0}$

(e):  $\lim_{x \rightarrow 2} \frac{x + 7}{(x - 2)^2}$  is  $\frac{7}{0}$ , so we check both sides:

LHL =  $\lim_{x \rightarrow 2^-} \frac{x + 7}{(x - 2)^2} = \frac{9}{(0^-)^2} = +\infty$

RHL =  $\lim_{x \rightarrow 2^+} \frac{x + 7}{(x - 2)^2} = \frac{9}{(0^+)^2} = +\infty$

LHL = RHL, so the original limit diverges to  $+\infty$


(f):  $\lim_{x \rightarrow -1} \frac{H(x + 1) - H(-1 - x)}{x + 1} = \lim_{x \rightarrow -1} \frac{\sqrt{(x + 1) + 2} - \sqrt{(-1 - x) + 2}}{x + 1}$   
 $= \lim_{x \rightarrow -1} \frac{\sqrt{(x + 1) + 2} - \sqrt{(-1 - x) + 2}}{x + 1} \cdot \left( \frac{\sqrt{(x + 1) + 2} + \sqrt{(-1 - x) + 2}}{\sqrt{(x + 1) + 2} + \sqrt{(-1 - x) + 2}} \right)$   
 $= \lim_{x \rightarrow -1} \frac{(x + 3) - (1 - x)}{(x + 1)(\sqrt{x + 3} + \sqrt{1 - x})} = \lim_{x \rightarrow -1} \frac{2x + 2}{(x + 1)(\sqrt{x + 3} + \sqrt{1 - x})}$   
 $= \lim_{x \rightarrow -1} \frac{2(x + 1)}{(x + 1)(\sqrt{x + 3} + \sqrt{1 - x})} = \lim_{x \rightarrow -1} \frac{2}{\sqrt{x + 3} + \sqrt{1 - x}} \stackrel{\text{DSP}}{=} \frac{2}{\sqrt{2} + \sqrt{2}} = \boxed{\frac{1}{\sqrt{2}}}$

**2.** [13 Points] Use translation, etc. to graph the following two functions:


$$f(x) = 5 + \sqrt{6 - x}$$

$$g(x) = \frac{1}{10}(x + 2)^2$$

**Solutions.** (a): This is  $y = \sqrt{x}$  translated **left by 6** to make  $y = \sqrt{x + 6}$  and *then reflected left-to-right* to make  $y = \sqrt{6 - x}$ , and finally translated **up by 5**. So it looks like:



(b): This is  $y = x^2$  translated **left by 2** and then **compressed vertically by 10**. So it looks like:



**3.** [15 Points] Suppose that  $f(x) = \frac{x + 7}{x - 3}$ . Compute  $f'(x)$  using the **limit definition of the derivative**.

$$\begin{aligned} \text{Solution. } f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{(x+h)+7}{(x+h)-3} - \frac{x+7}{x-3}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{x+h+7}{x+h-3} - \frac{x+7}{x-3}}{h} \cdot \left( \frac{(x+h-3)(x-3)}{(x+h-3)(x-3)} \right) = \lim_{h \rightarrow 0} \frac{(x+h+7)(x-3) - (x+7)(x+h-3)}{h(x+h-3)(x-3)} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + xh + 7x - 3x - 3h - 21 - (x^2 + xh - 3x + 7x + 7h - 21)}{h(x+h-3)(x-3)} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + xh + 7x - 3x - 3h - 21 - x^2 - xh + 3x - 7x - 7h + 21}{h(x+h-3)(x-3)} = \lim_{h \rightarrow 0} \frac{-10h}{h(x+h-3)(x-3)} \\ &= \lim_{h \rightarrow 0} \frac{-10}{(x+h-3)(x-3)} \stackrel{\text{DSP}}{=} \frac{-10}{(x-3)(x-3)} = \boxed{\frac{-10}{(x-3)^2}} \end{aligned}$$

**4.** [10 Points] Suppose that  $f(x) = x^2 - 7x - 12$ . Write the **equation of the tangent line** to the curve  $y = f(x)$  when  $x = -2$ . **\*\*Use the limit definition of the derivative when computing the derivative.\*\***

$$\text{Solution. } f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 7(x+h) - 12 - (x^2 - 7x - 12)}{h}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 7x - 7h - 12 - x^2 + 7x + 12}{h} = \lim_{h \rightarrow 0} \frac{2xh + h^2 - 7h}{h} \\
&= \lim_{h \rightarrow 0} 2x + h - 7 \stackrel{\text{DSP}}{=} 2x - 7.
\end{aligned}$$

Thus, the slope of the tangent line is  $f'(-2) = 2(-2) - 7 = -11$ .

We also have  $f(-2) = (-2)^2 - 7(-2) - 12 = 4 + 14 - 12 = 6$ , so the point is  $(-2, 6)$ . Therefore, the tangent line is

$$y - 6 = -11(x - (-2)), \quad \text{i.e.,} \quad \boxed{y = -11x - 16}$$

**5.** [12 Points] Suppose that  $f$  and  $g$  are functions, **and**

$$\bullet \lim_{x \rightarrow 7} f(x) = 5 \qquad \bullet \lim_{x \rightarrow 7} g(x) = -3 \qquad \bullet f(5) = 7 \qquad \bullet g(7) = \lim_{x \rightarrow 7} g(x).$$

Evaluate the following quantities and fully **justify** your answers. Do not just put down a value:

(a)  $\lim_{x \rightarrow 7} \sqrt{3f(x) - 7g(x)} =$

(b)  $\lim_{x \rightarrow 7} \frac{f(x)}{1 - x} =$

(c)  $g \circ f(5) =$

**Solutions.** (a): By the limit laws,  $\lim_{x \rightarrow 7} \sqrt{3f(x) - 7g(x)} = \sqrt{\lim_{x \rightarrow 7} 3f(x) - 7g(x)}$   
 $= \sqrt{3 \lim_{x \rightarrow 7} f(x) - 7 \lim_{x \rightarrow 7} g(x)} = \sqrt{3(5) - 7(-3)} = \sqrt{15 + 21} = \sqrt{36} = \boxed{6}$

(b): By the limit laws,  $\lim_{x \rightarrow 7} \frac{f(x)}{1 - x} = \frac{\lim_{x \rightarrow 7} f(x)}{\lim_{x \rightarrow 7} 1 - x} = \frac{5}{1 - 7} = \boxed{-\frac{5}{6}}$

(d): By the stated facts:  $g \circ f(5) = g(f(5)) = g(7) = \lim_{x \rightarrow 7} g(x) = \boxed{-3}$

**6.** [20 Points] Consider the function defined by

$$f(x) = \begin{cases} \sqrt{x-3} & \text{if } x > 3 \\ 1 & \text{if } x = 3 \\ 6 - 2x & \text{if } 0 < x < 3 \\ 16 - x^2 & \text{if } -4 < x \leq 0 \\ \frac{1}{x+4} & \text{if } x < -4 \end{cases}$$


(a) Carefully sketch the graph of  $f(x)$ .

(b) State the **Domain** of the function  $f(x)$ .

(c) Compute  $\begin{cases} \lim_{x \rightarrow 0^+} f(x) = \\ \lim_{x \rightarrow 0^-} f(x) = \\ \lim_{x \rightarrow 0} f(x) = \end{cases}$       (d) Compute  $\begin{cases} \lim_{x \rightarrow 3^+} f(x) = \\ \lim_{x \rightarrow 3^-} f(x) = \\ \lim_{x \rightarrow 3} f(x) = \end{cases}$

$$(e) \text{ Compute } \begin{cases} \lim_{x \rightarrow -4^+} f(x) = \\ \lim_{x \rightarrow -4^-} f(x) = \\ \lim_{x \rightarrow -4} f(x) = \end{cases}$$

**Solutions.** (a) Using translation/scaling for the various pieces, and then putting them together, here's the graph (axes not to scale, to fit better on the page)



(b): each of the functions is defined on the portion assigned to it, but  $x = -4$  is not in any of the portions. So  $\boxed{\text{the domain is all real numbers except } -4}$ , or if you prefer,  $\{x|x \neq -4\}$

$$(c): \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} 6 - 2x \stackrel{\text{DSP}}{=} \boxed{6}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} 16 - x^2 \stackrel{\text{DSP}}{=} \boxed{16}$$

$$\lim_{x \rightarrow 0} f(x) \boxed{\text{DNE}} \text{ because RHL} \neq \text{LHL}$$

$$(d): \lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} \sqrt{x-3} \stackrel{\text{DSP}}{=} \boxed{0}$$

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} 6 - 2x \stackrel{\text{DSP}}{=} \boxed{0}$$

$$\lim_{x \rightarrow 3} f(x) = \boxed{0} \text{ because RHL} = \text{LHL} = 0$$

$$(e): \lim_{x \rightarrow -4^+} f(x) = \lim_{x \rightarrow -4^+} 16 - x^2 \stackrel{\text{DSP}}{=} 16 - 16 = \boxed{0}$$

$$\lim_{x \rightarrow -4^-} f(x) = \lim_{x \rightarrow -4^-} \frac{1}{x+4} = \frac{1}{0^-} = \boxed{-\infty}$$

$$\lim_{x \rightarrow -4} f(x) \boxed{\text{DNE}} \text{ because RHL} \neq \text{LHL}$$