


### Solutions to Practice Test A for Midterm Exam 3

1. **(20 points)** You need to construct a box with a square base with a fixed volume of 24 cubic feet. The material for the bottom and top costs \$3 per square foot, and the material for the sides costs \$1 per square foot. What are the **dimensions** that minimize the cost required to build such a box? What is that **minimum cost**?

**Solution.** Here's the diagram:



The area of the base is  $x^2$ , so the base costs  $\$3x^2$ .

Similarly, the top also costs  $\$3x^2$ . So the base and top together cost  $\$6x^2$ .

Each of the four sides has area  $xy$ , and so costs  $\$xy$ . Together, the four sides cost  $\$4xy$ .

So the **total cost of the box (in dollars)** is  $6x^2 + 4xy$

The **volume** of the box is  $x^2y$ , which we set equal to 24. Solving for  $y$  gives  $y = 24x^{-2}$ .

So the cost of the box is  $C(x) = 6x^2 + 96x^{-1}$ .

The common sense bounds say  $x > 0$  [note that  $x = 0$  is impossible because  $x^2y = 24$ ]. [And  $y > 0$  then gives  $24/x^2 > 0$ , which tells us nothing new.]

So we must minimize  $C(x) = 6x^2 + 96x^{-1}$  on the domain  $(0, \infty)$ .

Differentiate:  $C'(x) = 12x - 96x^{-2}$ , which is **defined everywhere** on  $(0, \infty)$ .

Solving  $C' = 0$  gives  $12x^3 = 96$ , so  $x^3 = 8$ , so  $x = 2$  is the only critical number.

Using FDTAE, since  $C'(x) = 12x^{-2}(x^3 - 8)$ , our  $C'$  chart is:

$x$	$(0, 2)$	$(2, \infty)$
$C'(x)$	$-$	$+$
$C(x)$	$\searrow$	$\nearrow$

So by FDTAE,  $C$  has an absolute minimum at  $x = 2$  ft.

That gives  $y = 24/2^2 = 6$  ft and  $C(2) = 6(4) + 96/2 = \$72$ .

So the best box is  $\boxed{2 \text{ ft} \times 2 \text{ ft} \times 3 \text{ ft}}$  and  $\boxed{\text{costs } \$72}$

2. **(25 points)** Compute  $\int_1^3 x^2 - 3x \, dx$  using each of the following **two** different methods:

(a) The Fundamental Theorem of Calculus.

(b) Riemann Sums and the limit definition of the definite integral.

**Solution.** (a): Let  $f(x) = x^2 - 3x$ , and chop the interval  $[1, 3]$  into  $n$  equal-length intervals.

We have  $\Delta x = \frac{3-1}{n} = \frac{2}{n}$ , and so  $x_i = 1 + \frac{2i}{n}$ .

The  $n$ -th right-hand Riemann sum is therefore

$$R_n = \sum_{i=1}^n f(x_i)\Delta x = \sum_{i=1}^n f\left(1 + \frac{2i}{n}\right) \cdot \frac{2}{n} = \sum_{i=1}^n \left[ \left(1 + \frac{2i}{n}\right)^2 - 3\left(1 + \frac{2i}{n}\right) \right] \frac{2}{n}$$

$$\begin{aligned}
&= \sum_{i=1}^n \left[ 1 + \frac{4i}{n} + \frac{4i^2}{n^2} - 3 - \frac{6i}{n} \right] \frac{2}{n} = \sum_{i=1}^n \left[ -2 - \frac{2i}{n} + \frac{4i^2}{n^2} \right] \frac{2}{n} \\
&= -\frac{4}{n} \sum_{i=1}^n 1 - \frac{4}{n^2} \sum_{i=1}^n i + \frac{8}{n^3} \sum_{i=1}^n i^2 = -\frac{4}{n} \cdot n - \frac{4}{n^2} \cdot \frac{n(n+1)}{2} + \frac{8}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} \\
&= -4 - 2 \left( 1 + \frac{1}{n} \right) + \frac{4}{3} \left( 1 + \frac{1}{n} \right) \left( 2 + \frac{1}{n} \right)
\end{aligned}$$

Thus,  $\int_1^3 x^2 - 3x \, dx = \lim_{n \rightarrow \infty} R_n = -4 - 2(1) + \frac{4}{3}(1)(2) = -6 + \frac{8}{3} = \boxed{-\frac{10}{3}}$

(b):  $\int_1^3 x^2 - 3x \, dx = \left. \frac{1}{3}x^3 - \frac{3}{2}x^2 \right|_1^3 = \left( \frac{1}{3} \cdot 3^3 - \frac{3}{2} \cdot 3^2 \right) - \left( \frac{1}{3} \cdot 1^3 - \frac{3}{2} \cdot 1^2 \right)$   
 $= 9 - \frac{27}{2} - \frac{1}{3} + \frac{3}{2} = 9 - 12 - \frac{1}{3} = -3 - \frac{1}{3} = \boxed{-\frac{10}{3}}$

3. (30 points) Let  $f(x) = \frac{-x^2 + x + 2}{x^2 - 2x + 1} = \frac{-x^2 + x + 2}{(x-1)^2}$ .

For this function, discuss domain, vertical and horizontal asymptote(s), interval(s) of increase or decrease, local extreme value(s), concavity, and inflection point(s). Then use this information to present a detailed and labelled sketch of the curve  $y = f(x)$ .

**You may take my word for it that:**

$$f'(x) = \frac{x-5}{(x-1)^3} \quad \text{and} \quad f''(x) = \frac{-2x+14}{(x-1)^4}$$

**Solution.**

- Domain:  $f(x)$  has domain  $\{x|x \neq 1\}$
- VA: Vertical asymptote at  $x = 1$  (because denom= 0 there, and nowhere else).
- HA: Horizontal asymptote is  $y = -1$  on both sides since  $\lim_{x \rightarrow \pm\infty} f(x) = -1$  because

$$\lim_{x \rightarrow \pm\infty} \frac{-x^2 + x + 2}{x^2 - 2x + 1} \cdot \frac{\left(\frac{1}{x^2}\right)}{\left(\frac{1}{x^2}\right)} = \lim_{x \rightarrow \pm\infty} \frac{-1 + \frac{1}{x} + \frac{2}{x^2}}{1 - \frac{2}{x} + \frac{1}{x^2}} = -1$$

- First Derivative Information:

We know  $f'(x) = \frac{x-5}{(x-1)^3}$ . The critical points occur where  $f'$  is undefined or zero. The former happens when  $x = 1$ , but  $x = 1$  was not in the domain of the original function, so it isn't technically a critical number. The latter happens when  $x = 5$ . As a result,  $x = 5$  is the critical number. Our chart for  $f'$  is:

$x$	$(-\infty, 1)$	$(1, 5)$	$(5, \infty)$
$f'(x)$	+	-	+
$f(x)$	$\nearrow$	$\searrow$	$\nearrow$

So  $f$  is decreasing on  $(1, 5)$  and increasing on  $(-\infty, 1)$  and  $(5, \infty)$   
with a local minimum at  $x = 5$

(Remember that  $x = 1$  is a vertical asymptote, not a max.)

- Second Derivative Information:

Meanwhile,  $f'' = \frac{-2x+14}{(x-1)^4}$ , which is **always defined** away from the asymptote  $x = 1$ .

We have  $f'' = 0$  when  $x = 7$ . Thus, our  $f''$  chart is

$x$	$(-\infty, 1)$	$(1, 7)$	$(7, \infty)$
$f''(x)$	$+$	$+$	$-$
$f(x)$	$\cup$	$\cup$	$\cap$

So  $f$  is concave down on  $(7, \infty)$  and concave up on  $(-\infty, 1)$  and  $(1, 7)$

with an inflection point at  $x = 7$

[For graph, see separate PDF.]

4. (10 points) Compute the following limits.

$$(a) \lim_{x \rightarrow \infty} \frac{x^2 - x + 5}{3x^7 + x^6 - 2022}$$

$$(b) \lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 + 5x}}{6x + 7}$$

**Solution.** (a)  $\lim_{x \rightarrow \infty} \frac{x^2 - x + 5}{3x^7 + x^6 - 2022} \cdot \frac{1}{x^7} = \lim_{x \rightarrow \infty} \frac{x^{-5} - x^{-6} + 5x^{-7}}{3 + x^{-1} - 2022x^{-7}} = \frac{0}{3} = \boxed{0}$

(b)  $\lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 + 5x}}{6x + 7} \cdot \frac{1}{x} = \lim_{x \rightarrow -\infty} \frac{\sqrt{4x^2 + 5x} \cdot (-\sqrt{x^{-2}})}{6 + 7x^{-1}} = \lim_{x \rightarrow -\infty} \frac{-\sqrt{4 + 5x^{-1}}}{6 + 7x^{-1}} = \frac{-\sqrt{4}}{6} = \boxed{-\frac{1}{3}}$

5. (15 points) Compute the following definite and indefinite integrals.

$$(a) \int \frac{y^2 - 4y + 1}{\sqrt{y}} dy$$

$$(b) \int_{-\pi/4}^{5\pi/6} 3x + \cos x dx$$

**Solution.** (a)  $\int \frac{y^2 - 4y + 1}{\sqrt{y}} dy = \int y^{3/2} - 4y^{1/2} + y^{-1/2} dy = \boxed{\frac{2}{5}y^{5/2} - \frac{8}{3}y^{3/2} + 2y^{1/2} + C}$

(b)  $\int_{-\pi/4}^{5\pi/6} 3x + \cos x dx = \frac{3}{2}x^2 + \sin x \Big|_{-\pi/4}^{5\pi/6} = \left( \frac{3}{2} \left( \frac{5\pi}{6} \right)^2 + \sin \frac{5\pi}{6} \right) - \left( \frac{3}{2} \left( -\frac{\pi}{4} \right)^2 + \sin \left( -\frac{\pi}{4} \right) \right)$

$$= \frac{35^2\pi^2}{2 \cdot 6^2} + \frac{1}{2} - \frac{3\pi^2}{2 \cdot 4^2} - \left( -\frac{\sqrt{2}}{2} \right) = \boxed{\left( \frac{25}{24} - \frac{3}{32} \right) \pi^2 + \frac{1 + \sqrt{2}}{2}}$$